

















-  **7:** Collins LV, Eriksson K, Ulrich RG, Tarkowski A. Related Articles, Links
 Mucosal tolerance to a bacterial superantigen indicates a novel pathway to prevent toxic shock.
Infect Immun. 2002 May;70(5):2282-7.
PMID: 11953361 [PubMed - indexed for MEDLINE]
-  **8:** Murphey ED, Traber DL. Related Articles, Links
 Protective effect of tumor necrosis factor-alpha against subsequent endotoxemia in mice is mediated, in part, by interleukin-10.
Crit Care Med. 2001 Sep;29(9):1761-6.
PMID: 11546981 [PubMed - indexed for MEDLINE]
-  **9:** Le Roy D, Di Padova F, Adachi Y, Glauser MP, Calandra T, Heumann D. Related Articles, Links
 Critical role of lipopolysaccharide-binding protein and CD14 in immune responses against gram-negative bacteria.
J Immunol. 2001 Sep 1;167(5):2759-65.
PMID: 11509620 [PubMed - indexed for MEDLINE]
-  **10:** Singh BR, Sharma VD. Related Articles, Links
 Isolation and characterization of Salmonella Gallinarum cytotoxic factors.
Indian J Exp Biol. 2000 Nov;38(11):1152-8.
PMID: 11395962 [PubMed - indexed for MEDLINE]
-  **11:** Dinges MM, Schlievert PM. Related Articles, Links
 Role of T cells and gamma interferon during induction of hypersensitivity to lipopolysaccharide by toxic shock syndrome toxin 1 in mice.
Infect Immun. 2001 Mar;69(3):1256-64.
PMID: 11179286 [PubMed - indexed for MEDLINE]
-  **12:** Faure G, Villela C, Perales J, Bon C. Related Articles, Links
 Interaction of the neurotoxic and nontoxic secretory phospholipases A2 with the crotoxin inhibitor from Crotalus serum.
Eur J Biochem. 2000 Aug;267(15):4799-808.
PMID: 10903514 [PubMed - indexed for MEDLINE]
-  **13:** Dankesreiter S, Hoess A, Schneider-Mergener J, Wagner H, Miethke T. Related Articles, Links
 Synthetic endotoxin-binding peptides block endotoxin-triggered TNF-alpha production by macrophages in vitro and in vivo and prevent endotoxin-mediated toxic shock.
J Immunol. 2000 May 1;164(9):4804-11.
PMID: 10779788 [PubMed - indexed for MEDLINE]
-  **14:** Liu J, Ginis I, Spatz M, Hallenbeck JM. Related Articles, Links
 Hypoxic preconditioning protects cultured neurons against hypoxic stress via TNF-alpha and ceramide.
Am J Physiol Cell Physiol. 2000 Jan;278(1):C144-53.
PMID: 10644522 [PubMed - indexed for MEDLINE]

☐ **15:** [Denis-Mize KS, Price BM, Baker NR, Galloway DR.](#) [Related Articles, Links](#)



Analysis of immunization with DNA encoding *Pseudomonas aeruginosa* exotoxin A.

FEMS Immunol Med Microbiol. 2000 Feb;27(2):147-54.
PMID: 10640610 [PubMed - indexed for MEDLINE]

☐ **16:** [Heremans H, Dillen C, Groenen M, Matthys P, Billiau A.](#) [Related Articles, Links](#)



Role of interferon-gamma and nitric oxide in pulmonary edema and death induced by lipopolysaccharide.

Am J Respir Crit Care Med. 2000 Jan;161(1):110-7.
PMID: 10619806 [PubMed - indexed for MEDLINE]

☐ **17:** [Santanirand P, Harley VS, Dance DA, Drasar BS, Bancroft GJ.](#) [Related Articles, Links](#)



Obligatory role of gamma interferon for host survival in a murine model of infection with *Burkholderia pseudomallei*.

Infect Immun. 1999 Jul;67(7):3593-600.
PMID: 10377144 [PubMed - indexed for MEDLINE]

☐ **18:** [D'Souza MJ, Oettinger CW, Milton GV.](#) [Related Articles, Links](#)



Evaluation of microspheres containing cytokine neutralizing antibodies in endotoxemia.

Drug Dev Ind Pharm. 1999 Jun;25(6):727-34.
PMID: 10349558 [PubMed - indexed for MEDLINE]

☐ **19:** [de Bont N, Netea MG, Demacker PN, Verschueren I, Kullberg BJ, van Dijk KW, van der Meer JW, Stalenhoef AF.](#) [Related Articles, Links](#)



Apolipoprotein E knock-out mice are highly susceptible to endotoxemia and *Klebsiella pneumoniae* infection.

J Lipid Res. 1999 Apr;40(4):680-5.
PMID: 10191292 [PubMed - indexed for MEDLINE]

☐ **20:** [Lee FY, Li Y, Zhu H, Yang S, Lin HZ, Trush M, Diehl AM.](#) [Related Articles, Links](#)



Tumor necrosis factor increases mitochondrial oxidant production and induces expression of uncoupling protein-2 in the regenerating mice [correction of rat] liver.

Hepatology. 1999 Mar;29(3):677-87.
PMID: 10051468 [PubMed - indexed for MEDLINE]

Display	Summary	<input type="checkbox"/>	Show: 20	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Send to	Text	<input type="checkbox"/>
Items 1-20 of 80							Page	2	of 4 Next

[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act | Disclaimer](#)

Jun 12 2003 10:19:17